

THE LIFTING PROBLEM WITH THE FULL IDEAL

S. SHELAH

Abstract. We show that there are a cardinal μ , a σ -ideal $I \subseteq \mathcal{P}(\mu)$ and a σ -subalgebra $\mathcal{B}$ of subsets of μ extending I such that $\mathcal{B}/I$ satisfies the c.c.c. but the quotient algebra $\mathcal{B}/I$ has no lifting.

0. Introduction. In the present paper we prove the following theorem.

Theorem 0.1. *For some μ (in fact, $\mu = (2^{\aleph_0})^{++}$ suffices) there is a σ -ideal I on $\mathcal{P}(\mu)$ and a σ -subalgebra $\mathcal{B}$ of $\mathcal{P}(\mu)$ extending I such that $\mathcal{B}/I$ satisfies the c.c.c. but $\mathcal{B}/I$ has no lifting.*

This result answers a question of David Fremlin (see chapter on measure algebras in Fremlin [2]). Moreover, it solves the problem of topologizing a Category Base (see Detlefsen Szymański [3], Morgan [6], Shilling [11] and Szymański [12]).

Note that it is well known (Mokobodzki's theorem; see Fremlin [2]) that under CH, if $|\mathcal{B}/I| \leq (2^{\aleph_0})^+$ then this is impossible; i.e. the quotient algebra $\mathcal{B}/I$ has a lifting.

Toward the end we deal with having better μ .

I thank Andrzej Szymański for asking me the question and Max Burke and Mariusz Rabus for corrections.

Notation: Our notation is rather standard. All cardinals are assumed to be infinite and usually they are denoted by λ , κ , μ .

In Boolean algebras we use $\cap$ (and $\cap$), $\cup$ (and $\cup$) and $-$ for the Boolean operations.

Date: February 1, 2008.

1991 *Mathematics Subject Classification.* Primary: 03E05, 28A51.

I would like to thank Alice Leonhardt for the beautiful typing.

The research was partially supported by "Basic Research Foundation" of the Israel Academy of Sciences and Humanities. Publication 636.

1. The proof of Theorem 0.1.

Main Lemma 1.1. *Suppose that*

- (a) μ, λ are cardinals satisfying $\mu = \mu^{\aleph_0}, \lambda \leq 2^\mu$,
- (b) $\mathfrak{B}$ is a complete c.c.c. Boolean algebra,
- (c) $x_i \in \mathfrak{B} \setminus \{0\}$ for $i < \lambda$,
- (d) for each sequence $\langle (u_i, f_i) : i < \lambda \rangle$ such that $u_i \in [\lambda]^{\leq \aleph_0}, f_i \in {}^{u_i}2$ there are $n < \omega$ (but $n > 0$) and $i_0 < i_1 < \dots < i_{n-1}$ in λ such that:
 - (α) the functions $f_{i_0}, \dots, f_{i_{n-1}}$ are compatible,
 - (β) $\mathfrak{B} \models \bigcap_{\ell < n} x_{i_\ell} = 0$.

Then

- ($\oplus$) there are a σ -ideal I on $\mathcal{P}(\mu)$ and a σ -algebra $\mathfrak{A}$ of subsets of μ extending I such that $\mathfrak{A}/I$ satisfies the c.c.c. and the natural homomorphism $\mathfrak{A} \longrightarrow \mathfrak{A}/I$ cannot be lifted.

PROOF Without loss of generality the algebra $\mathfrak{B}$ has cardinality $\lambda^{\aleph_0}$ ($\leq 2^\mu$). Let $\langle Y_b : b \in \mathfrak{B} \rangle$ be a sequence of subsets of μ such that any non-trivial countable Boolean combination of the Y_b 's is non-empty (possible by [1] as $\mu = \mu^{\aleph_0}$ and the algebra $\mathfrak{B}$ has cardinality $\leq 2^\mu$; see background in [4]). Let $\mathfrak{A}_0$ be the Boolean subalgebra of $\mathcal{P}(\mu)$ generated by $\{Y_b : b \in \mathfrak{B}\}$. So $\{Y_b : b \in \mathfrak{B}\}$ freely generates $\mathfrak{A}_0$ and hence there is a unique homomorphism h_0 from $\mathfrak{A}_0$ into $\mathfrak{B}$ satisfying $h_0(Y_b) = b$.

A Boolean term σ is hereditarily countable if σ belongs to the closure Σ of the set of terms $\bigcap_{i < i^*} y_i$ for $i^* < \omega_1$ under composition and under $-y$.

Let $\mathcal{E}$ be the set of all equations $\mathbf{e}$ of the form $0 = \sigma(b_0, b_1, \dots, b_n, \dots)_{n < \omega}$ which hold in $\mathfrak{B}$, where σ is hereditarily countable. For $\mathbf{e} \in \mathcal{E}$ let $\text{cont}(\mathbf{e})$ be the set of $b \in \mathfrak{B}$ mentioned in it (i.e. $\{b_n : n < \omega\}$) and let $Z_{\mathbf{e}} \subseteq \mu$ be the set $\sigma(Y_{b_0}, Y_{b_1}, \dots, Y_{b_n}, \dots)_{n < \omega}$.

Let I be the σ -ideal of $\mathcal{P}(\mu)$ generated by the family $\{Z_{\mathbf{e}} : \mathbf{e} \in \mathcal{E}\}$ and let $\mathfrak{A}_1$ be the Boolean Algebra of subsets of $\mathcal{P}(\mu)$ generated by $I \cup \{Y_b : b \in \mathfrak{B}\}$.

Claim 1.1.1. $I \cap \mathfrak{A}_0 = \text{Ker}(h_0)$.

Proof of the claim: Plainly $\text{Ker}(h_0) \subseteq I \cap \mathfrak{A}_0$. For the converse inclusion it is enough to consider elements of $\mathfrak{A}_0$ of the form

$$Y = \bigcap_{\ell=1}^n Y_{b_\ell} - \bigcup_{\ell=n+1}^{2n} Y_{b_\ell}.$$

If $\mathfrak{B} \models \text{"}\bigcap_{\ell=1}^n b_\ell - \bigcup_{\ell=n+1}^{2n} b_\ell = 0\text{"}$ then easily $h_0(Y) = 0$. So assume that

$$\mathfrak{B} \models \text{"}c = \bigcap_{\ell=1}^n b_\ell - \bigcup_{\ell=n+1}^{2n} b_\ell \neq 0\text{"},$$

and we shall prove $Y \notin I$. Let $Z \in I$, so for some $\mathbf{e}_m \in \mathcal{E}$ for $m < \omega$ we have $Z \subseteq \bigcup_{m < \omega} Z_{\mathbf{e}_m}$. Let g be a homomorphism from $\mathfrak{B}$ into the 2-element Boolean Algebra $\mathfrak{B}_0 = \{0, 1\}$ such that $g(c) = 1$, and g respects all the equations $\mathbf{e}_m$ (including those of the form $b = \bigcup_{k < \omega} b_k$; possible by the Sikorski theorem).

By the choice of the Y_b 's, there is $\alpha < \mu$ such that:

if $b \in \{b_\ell : \ell = 1, \dots, 2n\} \cup \bigcup_{m < \omega} \text{cont}(\mathbf{e}_m)$ then

$$g(b) = 1 \Leftrightarrow \alpha \in Y_b.$$

So easily $\alpha \notin Z_{\mathbf{e}_m}$ for $m < \omega$, and $\alpha \in \bigcap_{\ell=1}^n Y_{b_\ell} \setminus \bigcup_{\ell=n+1}^{2n} Y_{b_\ell}$, so Y is not a subset of Z . As Z was an arbitrary element of I we get $Y \notin I$, so we have finished proving 1.1.1.

It follows from 1.1.1 that we can extend h_0 (the homomorphism from $\mathfrak{A}_0$ onto $\mathfrak{B}$) to a homomorphism h_1 from $\mathfrak{A}_1$ onto $\mathfrak{B}$ with $I = \text{Ker}(h_1)$. Let $\mathfrak{A}_2$ be the σ -algebra of subsets of μ generated by $\mathfrak{A}_1$.

Claim 1.1.2. *For every $Y \in \mathfrak{A}_2$ there is $b \in \mathfrak{B}$ such that $Y \equiv Y_b \pmod{I}$. Consequently, $\mathfrak{A}_2 = \mathfrak{A}_1$.*

Proof of the claim: Let $Y \in \mathfrak{A}_2$. Then Y is a (hereditarily countable) Boolean combination of some Y_{b_ℓ} ($\ell < \omega$) and Z_n ($n < \omega$), where $b_\ell \in \mathfrak{B}$, $Z_n \in I$. Let $Z_n \subseteq \bigcup_{m < \omega} Z_{\mathbf{e}_{n,m}}$, where $\mathbf{e}_{n,m} \in \mathcal{E}$, and say

$$Y = \sigma(Y_{b_0}, Z_0, Y_{b_1}, Z_1, \dots, Y_{b_n}, Z_n, \dots)_{n < \omega}.$$

Let $\mathbf{e}_{n,m}$ be $0 = \sigma_{n,m}(b_{n,m,0}, b_{n,m,1}, \dots)$. Then clearly $\bigcup_{n,m < \omega} Z_{\mathbf{e}_{n,m}} \in I$ (use the definition of I). In $\mathfrak{B}$, let $b = \sigma(b_0, 0, b_1, 0, \dots, b_n, 0, \dots)$ and let $\sigma^* = \sigma^*(b_0, b_1, \dots, b_{n,m,\ell}, \dots)_{n,m,\ell < \omega}$ be the following term

$$\begin{aligned} & \bigcup_{n,m} \sigma_{n,m}(b_{n,m,0}, b_{n,m,1}, \dots) \cup (b - \sigma(b_0, 0, b_1, 0, \dots, b_m, 0, \dots)) \cup \\ & \cup (\sigma(b_0, 0, b_1, 0, \dots, b_n, 0, \dots) - b) \cup 0. \end{aligned}$$

Clearly $\mathfrak{B} \models \text{"}0 = \sigma^*\text{"}$, so the equation $\mathbf{e}$ defined as $0 = \sigma^*$ belongs to $\mathcal{E}$, and thus $Z_{\mathbf{e}}$ is well defined. It follows from the definition of σ^* that $(Y \setminus Y_b) \cup (Y_b \setminus Y) \subseteq Z_{\mathbf{e}} \in I$.

So we can sum up:

- (a) I is an $\aleph_1$ -complete ideal of $\mathcal{P}(\mu)$,
- (b) $\mathfrak{A}_1$ is a σ -algebra of subsets of μ ,
- (c) $I \subseteq \mathfrak{A}_1$,
- (d) h_1 is a homomorphism from $\mathfrak{A}_1$ onto $\mathfrak{B}$, with kernel I ,
- (e) $\mathfrak{B}$ is a complete c.c.c. Boolean algebra.

This is exactly as required, so the “only” point left is

Claim 1.1.3. *The homomorphism h_1 cannot be lifted.*

Proof of the claim: Assume that h_1 can be lifted, so there is a homomorphism $g_1 : \mathfrak{B} \longrightarrow \mathfrak{A}_1$ such that $h_1 \circ g_1 = \text{id}_{\mathfrak{B}}$.

For $i < \lambda$ let $Z_i = (g_1(x_i) - Y_{x_i}) \cup (Y_{x_i} - g_1(x_i))$, so by the assumption on g_1 necessarily $Z_i \in I$. Consequently we can find $\mathbf{e}_{i,n} \in \mathcal{E}$ for $n < \omega$ such that $Z_i \subseteq \bigcup_{n < \omega} Z_{\mathbf{e}_{i,n}}$. Let $W_i = \{x_i\} \cup \bigcup_{n < \omega} \text{cont}(\mathbf{e}_{i,n})$, so $W_i \subseteq \mathfrak{B}$ is countable. Let $\mathfrak{B}'$ be the subalgebra of $\mathfrak{B}$ generated by $\bigcup_{i < \lambda} W_i$. Clearly $|\mathfrak{B}'| = \lambda$, so there

is a one-to-one function t from λ onto $\mathfrak{B}'$. Put $u_i = t^{-1}(W_i) \in [\lambda]^{\leq \aleph_0}$.

For each i there is a homomorphism f_i from $\mathfrak{B}$ into the 2-element Boolean Algebra $\{0, 1\}$ such that $f_i(x_i) = 1$ and f_i respects all the equations $\mathbf{e}_{i,n}$ for $n < \omega$ (as in the proof of 1.1.1). Let $f'_i : u_i \longrightarrow \{0, 1\}$ be defined by $f'_i(\alpha) = f_i(t(\alpha))$. Then by clause (d) of the hypothesis there are $n < \omega$ and $i_0 < \dots < i_{n-1} < \lambda$ such that:

- (α) the functions $f'_{i_0}, \dots, f'_{i_{n-1}}$ are compatible,
- (β) $\mathfrak{B} \models \bigcap_{\ell < n} x_{i_\ell} = 0$.

Hence

- (α') the functions $f_{i_0} \upharpoonright W_{i_0}, \dots, f_{i_{n-1}} \upharpoonright W_{i_{n-1}}$ are compatible¹, call their union g .

Now let $\alpha < \mu$ be such that:

$$(\otimes_1) \quad \ell < n \ \& \ b \in W_{i_\ell} \quad \Rightarrow \quad [\alpha \in Y_b \Leftrightarrow g(b) = 1]$$

(it exists by the choice of the Y_b 's and (α')).

By ($\otimes_1$) and the choice of f_{i_ℓ} we have:

$$(\otimes_2) \quad \alpha \in Y_{x_{i_\ell}}$$

(because $f_{i_\ell}(x_{i_\ell}) = 1$) and

$$(\otimes_3) \quad \alpha \notin Z_{\mathbf{e}_{i_\ell, n}} \text{ for } n < \omega$$

(because f_{i_ℓ} respects $\mathbf{e}_{i_\ell, n}$ and $\text{cont}(\mathbf{e}_{i_\ell, n}) \subseteq W_{i_\ell}$) and

$$(\otimes_4) \quad \alpha \notin Z_{i_\ell}$$

¹as functions, not as homomorphisms

(by $(\otimes_3)$ as $Z_{i_\ell} \subseteq \bigcup_{n < \omega} Z_{e_{i_\ell, n}}$).

So $\alpha \in Y_{x_{i_\ell}} \setminus Z_{i_\ell}$ and thus $\alpha \in g_1(x_{i_\ell})$. Hence $\alpha \in \bigcap_{\ell < n} g_1(x_{i_\ell})$. Since g_1 is a homomorphism we have

$$\bigcap_{\ell < n} g_1(x_{i_\ell}) = g_1\left(\bigcap_{\ell < n} x_{i_\ell}\right) = g_1(0) = \emptyset$$

(we use clause (β) above). A contradiction. ■_{1.1}

Remark 1.2. 1. Concerning the assumptions of 1.1, note that they seem closely related to

$(\oplus_\mu)$ there is a c.c.c. Boolean Algebra $\mathfrak{B}$ of cardinality $\leq \lambda$ which is not the union of $\leq \mu$ ultrafilters (i.e. $d(\mathfrak{B}) > \mu$).

(See the proof of 1.7 below).

2. Concerning $(\oplus_\mu)$, by [8], if $\lambda = \mu^+$, $\mu = \mu^{\aleph_0}$ then there is no such Boolean algebra. By [9], it is consistent then $\lambda = \mu^{++} \leq 2^\mu$, $\aleph_0 < \mu = \mu^{<\mu}$ and $(\oplus_\mu)$ above holds using (see below) a Boolean algebra of the form $BA(W)$, $W \subseteq [\lambda]^3$, $(\forall u_1 \neq u_2 \in W)(|u_1 \cap u_2| \leq 1)$. Hajnal, Juhasz and Szentmiklossy [5] prove the existence of a c.c.c. Boolean algebra $\mathfrak{B}$ with $d(\mathfrak{B}) = \mu$ of cardinality 2^μ when there is a Jonsson algebra on μ (or μ is a limit cardinal) using $BA(W)$, $W \subseteq [\lambda]^{<\aleph_0}$, $u \neq v \in W \Rightarrow |u \cap v| < |u|/2$. The claim we need is close to this. On the existence of Jonsson cardinals (and its history) see [10]. Of course, also in 1.7 if μ is not strong limit, instead “ M is a Jonsson algebra on μ ” it suffices that “ M is not the union of $< \mu$ subalgebras”. Rabus Shelah [7] prove the existence of a c.c.c. Boolean Algebra $\mathfrak{B}$ with $d(\mathfrak{B}) = \mu$ for every μ .

Definition 1.3. 1. For a set u let

$$\text{pfil}(u) \stackrel{\text{def}}{=} \{w : w \subseteq \mathcal{P}(u), u \in w, w \text{ is upward closed and} \\ \text{if } (u_1, u_2) \text{ is a partition of } u \text{ then } u_1 \in w \text{ or } u_2 \in w\}$$

[pfil stands for “pseudo-filter”].

2. The canonical (pfil) w of u for a finite set u is

$$\text{half}(u) = \{v \subseteq u : |v| \geq |u|/2\}.$$

3. We say that $(W, \mathbf{w})$ is a λ -candidate if:

- (a) $W \subseteq [\lambda]^{<\aleph_0}$,
- (b) $\mathbf{w}$ is a function with domain W ,
- (c) $\mathbf{w}(u) \in \text{pfil}(u)$ for $u \in W$
- (d) if $v \in [\lambda]^{<\aleph_0}$ then $\text{cl}_{(W, \mathbf{w})}(v) \stackrel{\text{def}}{=} \{u \in W : u \cap v \in \mathbf{w}(u)\}$ is finite.

4. We say W is a λ -candidate if $(W, \text{half} \restriction W)$ is a λ -candidate.
5. Instead of λ we can use any ordinal (or even set).
6. We say that $\mathcal{U} \subseteq \lambda$ is $(W, \mathbf{w})$ -closed if for each $u \in W$

$$u \cap \mathcal{U} \in \mathbf{w}(u) \quad \Rightarrow \quad u \subseteq \mathcal{U}.$$

Definition 1.4. 1. For a λ -candidate $(W, \mathbf{w})$ let $BA(W, \mathbf{w})$ be the Boolean algebra generated by $\{x_i : i < \lambda\}$ freely except

$$\bigcap_{i \in u} x_i = 0 \quad \text{for} \quad u \in W.$$

2. For a λ -candidate W , let

$$BA(W) = BA(W, \text{half} \restriction W).$$

3. For a λ -candidate $(W, \mathbf{w})$ let $BA^c(W, \mathbf{w})$ be the completion of $BA(W, \mathbf{w})$; similarly $BA^c(W)$.

Proposition 1.5. *Let $(W, \mathbf{w})$ be a λ -candidate. Then the Boolean algebra $BA(W, \mathbf{w})$ satisfies the c.c.c. and has cardinality λ , so $BA^c(W, \mathbf{w})$ satisfies the c.c.c. and has cardinality $\leq \lambda^{\aleph_0}$.*

PROOF Let $b_\alpha = \sigma_\alpha(x_{i_{\alpha,0}}, \dots, x_{i_{\alpha, n_\alpha-1}})$ be nonzero members of $BA(W, \mathbf{w})$ (for $\alpha < \omega_1$ and σ_α a Boolean term). Without loss of generality $\sigma_\alpha = \sigma$, $n_\alpha = n(*)$ and $i_{\alpha,0} < i_{\alpha,1} < \dots < i_{\alpha, n_\alpha-1}$, and $\langle \langle i_{\alpha, \ell} : \ell < n(*) \rangle : \alpha < \omega_1 \rangle$ forms a Δ -system, so

$$i_{\alpha_1, \ell_1} = i_{\alpha_2, \ell_2} \ \& \ \alpha_1 \neq \alpha_2 \quad \Rightarrow \quad \ell_1 = \ell_2 \ \& \ (\forall \alpha < \omega_1)(i_{\alpha, \ell_1} = i_{\alpha_1, \ell_1}).$$

Also we can replace b_α by any nonzero $b'_\alpha \leq b_\alpha$, so without loss of generality for some $s_\alpha \subseteq n(*)$ ($= \{0, \dots, n(*) - 1\}$) we have

$$b_\alpha = \bigcap_{\ell \in s_\alpha} x_{i_{\alpha, \ell}} \cap \bigcap_{\ell \in n(*) \setminus s_\alpha} (-x_{i_{\alpha, \ell}}) > 0$$

and without loss of generality $s_\alpha = s$. Put (for $\alpha < \omega_1$)

$$\mathbf{u}_\alpha \stackrel{\text{def}}{=} \{u \in W : u \cap \{i_{\alpha, \ell} : \ell \in s\} \in \mathbf{w}(u)\}$$

and note that these sets are finite (remember 1.3(3d)). Hence the sets

$$u_\alpha = \bigcup \{u : u \in \mathbf{u}_\alpha\}$$

are finite. Without loss of generality $\langle \{i_{\alpha, \ell} : \ell < n(*)\} \cup u_\alpha : \alpha < \omega_1 \rangle$ is a Δ -system. Now let $\alpha \neq \beta$ and assume $b_\alpha \cap b_\beta = 0$. Clearly we have

$$b_\alpha \cap b_\beta = \bigcap_{\ell \in s} (x_{i_{\alpha, \ell}} \cap x_{i_{\beta, \ell}}) \cap \bigcap_{\ell \in n(*) \setminus s} (-x_{i_{\alpha, \ell}} \cap -x_{i_{\beta, \ell}}).$$

Note that, by the Δ -system assumption, the sets $\{i_{\alpha,\ell}, i_{\beta,\ell} : \ell \in s\}$, $\{i_{\alpha,\ell}, i_{\beta,\ell} : \ell \in n(*) \setminus s\}$ are disjoint. So why is $b_\alpha \cap b_\beta$ zero? The only possible reason is that for some $u \in W$ we have $u \subseteq \{i_{\alpha,\ell}, i_{\beta,\ell} : \ell \in s\}$. Thus

$$u = (u \cap \{i_{\alpha,\ell} : \ell \in s\}) \cup \{u \cap \{i_{\beta,\ell} : \ell \in s\}\}$$

and without loss of generality $u \cap \{i_{\alpha,\ell} : \ell \in s\} \in \mathbf{w}(u)$. Hence $u \in \mathbf{u}_\alpha$ and therefore $u \subseteq u_\alpha$. Now we may easily finish the proof. $\blacksquare_{1.5}$

Remark 1.6. If we define a (λ, κ) -candidate weakening clause (d) to

$$(d)_\kappa \quad v \in [\lambda]^{<\aleph_0} \quad \Rightarrow \quad \kappa > |\{u \in W : u \cap v \in \mathbf{w}(u)\}|,$$

then the algebra $BA(W, \mathbf{w})$ satisfies the κ^+ -c.c.c.

[Why? We repeat the proof of Proposition 1.5 replacing $\aleph_1$ with κ . There is a difference only when $\mathbf{u}_\alpha$ has cardinality $< \kappa$ (instead being finite) and (being the union of $< \kappa$ finite sets) also u_α has cardinality $\mu_\alpha < \kappa$. Wlog $\mu_\alpha = \mu < \kappa$. Clearly the set

$$S \stackrel{\text{def}}{=} \{\delta < \kappa^+ : \text{cf}(\delta) = \mu^+\}$$

is a stationary subset of κ^+ , so for some stationary subset S^* of S and $\alpha(*) < \kappa$ we have:

$$(\forall \alpha \in S^*)(u_\alpha \cap \alpha \subseteq \alpha^* \quad \& \quad u_\alpha \subseteq \min(S^* \setminus (\alpha + 1))).$$

Let us define $u_\alpha^* = u_\alpha \cup \{i_{\alpha,\ell} : \ell \in s\} \setminus \alpha(*)$. Wlog $\langle u_\alpha^* : \alpha \in S^* \rangle$ is a Δ -system. The rest should be clear.]

Theorem 1.7. Assume that there is a Jonsson algebra on μ , $\lambda = 2^\mu$, and

$$(\forall \alpha < \mu)(|\alpha|^{\aleph_0} < \mu = \text{cf}(\mu)).$$

Then for some λ -candidate $(W, \mathbf{w})$ the Boolean algebra $BA^c(W, \mathbf{w})$ and λ satisfy the assumptions **(b)**–**(d)** of 1.1.

PROOF Let $F : [\mu]^{<\aleph_0} \longrightarrow \mu$ be such that

$$(\forall A \in [\mu]^\mu)[F''([A]^{<\aleph_0} \setminus [A]^{<2}) = \mu]$$

(well known and easily equivalent to the existence of a Jonsson algebra).

Let $\langle \bar{A}^\alpha : \alpha < 2^\mu \rangle$ list the sequences $\bar{A} = \langle A_i : i < \mu \rangle$ such that

- $A_i \in [2^\mu]^\mu$,
- $(\forall i < \mu)(\exists \alpha)(A_i \subseteq [\mu \times \alpha, \mu \times \alpha + \mu))$, and
- $i < j < \mu \quad \Rightarrow \quad A_i \cap A_j = \emptyset$.

Without loss of generality we have $A_i^\alpha \subseteq \mu \times (1 + \alpha)$ and each $\bar{A}$ is equal to $\bar{A}^\alpha$ for 2^μ ordinals α . Clearly $\text{otp}(A_i^\alpha) = \mu$.

By induction on $\alpha < 2^\mu$ we choose pairs $(W_\alpha, \mathbf{w}_\alpha)$ and functions F_α such that

- (α) $(W_\alpha, \mathbf{w}_\alpha)$ is a $\mu \times (1 + \alpha)$ -candidate,
- (β) $\beta < \alpha$ implies $W_\beta = W_\alpha \cap [\mu \times (1 + \beta)]^{<\aleph_0}$ and $\mathbf{w}_\beta = \mathbf{w}_\alpha \upharpoonright W_\beta$,
- (γ) F_α is a one-to-one function from the set

$$\{u : u \subseteq [\mu \times (1 + \alpha), \mu \times (1 + \alpha + 1)) \text{ finite with } \geq 2 \text{ elements}\}$$
 into $\bigcup_{i < \mu} A_i^\alpha$,
- (δ) $W_{\alpha+1} = W_\alpha \cup \{u \cup \{F_\alpha(u)\} : u \in W_\alpha^*\}$, where

$$W_\alpha^* = \{u : u \text{ a subset of } [\mu \times (1 + \alpha), \mu \times (1 + \alpha + 1)) \text{ such that } \aleph_0 > |u| \geq 2\},$$
- (ε) for any (finite) $u \in W_\alpha^*$ we have

$$\mathbf{w}_{\alpha+1}(u \cup \{F_\alpha(u)\}) = \{v \subseteq u \cup \{F_\alpha(u)\} : u \subseteq v \text{ or } F_\alpha(u) \in v \text{ \& } v \cap u \neq \emptyset\},$$
- (ζ) F_α is such that for any subset X of $J_\alpha = [\mu \times (1 + \alpha), \mu \times (1 + \alpha + 1))$ of cardinality μ and $i < \mu$ and $\gamma \in A_i^\alpha$ for some finite subset u of X with ≥ 2 elements we have $F_\alpha(u) \in A_i^\alpha \setminus \gamma$.

There is no problem to carry out the definition so that clauses (β)–(ζ) are satisfied (to define functions F_α use the function F chosen at the beginning of the proof). Then $(W_\alpha, \mathbf{w}_\alpha)$ is defined for each $\alpha < 2^\mu$ (at limit stages α we take $W_\alpha = \bigcup_{\beta < \alpha} W_\beta$, $\mathbf{w}_\alpha = \bigcup_{\beta < \alpha} \mathbf{w}_\beta$, of course).

Claim 1.7.1. *For each $\alpha < 2^\mu$, $(W_\alpha, \mathbf{w}_\alpha)$ is a $\mu \times (1 + \alpha)$ -candidate.*

Proof of the claim: We should check the requirements of 1.3(3). Clauses (a), (b) there are trivially satisfied. For the clause (c) note that every element u of W_α is of the form $u' \cup \{F_\beta(u')\}$ for some $\beta < \alpha$ and $u' \in W_\beta^*$. Now, if $u = u_0 \cup u_1$ then either one of u_0, u_1 contains u' or one of the two sets contains $F_\beta(u')$ and has non-empty intersection with u' . In both cases we are done. Regarding the demand (d) of 1.3(3), note that if

$$v \in [2^\mu]^{<\aleph_0}, \quad u \in W_\alpha, \quad u = u' \cup \{F_\beta(u')\}, \quad u' \in W_\beta^*, \quad \beta < \alpha$$

and $v \cap u \in \mathbf{w}_{\beta+1}(u)$ then $v \cap u' \neq \emptyset$ and either $u' \subseteq v$ or $F_\beta(u') \in u$. Hence, using the fact that the functions F_γ are one-to-one, we easily show that for every $v \in [2^\mu]^{<\aleph_0}$ the set

$$\{u \in W_\alpha : u \cap v \in \mathbf{w}_\alpha(u)\}$$

is finite (remember the definition of $\mathbf{w}_{\beta+1}$), finishing the proof of the claim.

Let $W = \bigcup_\alpha W_\alpha$, $\mathbf{w} = \bigcup_\alpha \mathbf{w}_\alpha$, $\mathfrak{B} = BA^c(W, \mathbf{w})$. It follows from 1.7.1 that $(W, \mathbf{w})$ is a λ -candidate. The main point of the proof of the theorem is clause (**d**) of the assumptions of 1.1. So let $f_\alpha : u_\alpha \longrightarrow \{0, 1\}$ for $\alpha < 2^\mu$, $u_\alpha \in [2^\mu]^{\leq \aleph_0}$, be given. For each $\alpha < 2^\mu$, by the assumption that

$(\forall \beta < \mu)[|\beta|^{\aleph_0} < \mu = \text{cf}(\mu)]$ and by the Δ -lemma, we can find $X_\alpha \in [\mu]^\mu$ such that $\langle f_{\mu \times \alpha + \zeta} : \zeta \in X_\alpha \rangle$ forms a Δ -system with heart f_α^* . Let

$G = \{g : g \text{ is a partial function from } 2^\mu \text{ to } \{0, 1\} \text{ with countable domain}\}.$

For each $g \in G$ let $\langle \gamma(g, i) : i < i(g) \rangle$ be a maximal sequence such that $g \subseteq f_{\gamma(g, i)}^*$ and

$$\text{Dom}(f_{\gamma(g, i)}^*) \cap \text{Dom}(f_{\gamma(g, j)}^*) = \text{Dom}(g) \quad \text{for } j < i$$

(just choose $\gamma(g, i)$ by induction on i).

By induction on $\zeta \leq \omega_1$, we choose $Y_\zeta, G_\zeta, Z_\zeta$ and $U_{\zeta, g}$ such that

- (a) $Y_\zeta \in [2^\mu]^{\leq \mu}$ is increasing continuous in ζ ,
- (b) $Z_\zeta \stackrel{\text{def}}{=} \bigcup \{\text{Dom}(f_\gamma) : (\exists \alpha \in Y_\zeta)[\mu \times \alpha \leq \gamma < \mu \times (\alpha + 1)]\},$
- (c) $G_\zeta = \{g \in G : \text{Dom}(g) \subseteq Z_\zeta\},$
- (d) for $g \in G_\zeta$ we have: $U_{\zeta, g}$ is $\{i : i < i(g)\}$ if $i(g) < \mu^+$ and otherwise it is a subset of $i(g)$ of cardinality μ such that

$$j \in U_{\zeta, g} \Rightarrow \text{Dom}(f_{\gamma(g, j)}^*) \cap Z_\zeta = \text{Dom}(g),$$

- (e) $Y_{\zeta+1} = Y_\zeta \cup \{\gamma(g, j) : g \in G_\zeta \text{ and } j \in U_{\zeta, g}\}.$

Let $Y = Y_{\omega_1}$. Let $\{(g_\varepsilon, \xi_\varepsilon) : \varepsilon < \varepsilon(*)\}$, $\varepsilon(*) \leq \mu$, list the set of pairs (g, ξ) such that $\xi < \omega_1$, $g \in G_\xi$ and $i(g) \geq \mu^+$. We can find $\langle \zeta_\varepsilon : \varepsilon < \varepsilon(*) \rangle$ such that $\langle \gamma(g_\varepsilon, \zeta_\varepsilon) : \varepsilon < \varepsilon(*) \rangle$ is without repetition and $\zeta_\varepsilon \in U_{g_\varepsilon, \xi_\varepsilon}$. Then for some $\alpha < 2^\mu \setminus Y_{\omega_1}$ we have

$$(\forall \varepsilon < \varepsilon(*))(A_\varepsilon^\alpha = \{\mu \times \gamma(g_\varepsilon, \zeta_\varepsilon) + \Upsilon : \Upsilon \in X_{\gamma(g_\varepsilon, \zeta_\varepsilon)}\}).$$

Now let $g = f_\alpha^* \upharpoonright Z_{\omega_1}$. Then for some $\zeta_0(*) < \omega_1$ we have $g \in G_{\zeta_0(*)}$ and thus $U_{g, \zeta} \subseteq i(g)$ for $\zeta \in [\zeta_0(*), \omega_1)$ and $\langle \gamma(g, i) : i < i(g) \rangle$ are well defined. Now, α exemplifies that $i(g) < \mu^+$ is impossible (see the maximality of $i(g)$, as otherwise $i < i(g) \Rightarrow \gamma(g, i) \in Y_{\zeta_0(*)+1} \subseteq Y_{\omega_1}$).

Next, for each $\gamma \in X_\alpha$, $\text{Dom}(f_{\mu \times \alpha + \gamma})$ is countable and hence for some $\zeta_{1, \gamma} < \omega_1$ we have $\text{Dom}(f_{\mu \times \alpha + \gamma}) \cap Z_{\omega_1} \subseteq Z_{\zeta_{1, \gamma}(*)}$. As $\text{cf}(\mu) > \aleph_1$ necessarily for some $\zeta_1(*) < \omega_1$ we have that $X'_\alpha \stackrel{\text{def}}{=} \{\gamma \in X_\alpha : \zeta_{1, \gamma} \leq \zeta_1(*)\} \in [\mu]^\mu$, and without loss of generality $\zeta_1(*) \geq \zeta_0(*)$.

So for some $\varepsilon < \varepsilon(*) \leq \mu$ we have $g_\varepsilon = g$ & $\xi_\varepsilon = \zeta_1(*) + 1$. Let $\Upsilon_\varepsilon = \gamma(g_\varepsilon, \zeta_\varepsilon)$. Clearly

- (*)₁ $f_\alpha^*, f_{\Upsilon_\varepsilon}^*$ are compatible (and countable),
- (*)₂ $\langle f_{\mu \times \alpha + \gamma} : \gamma \in X'_\alpha \rangle$ is a Δ -system with heart f_α^* .

So possibly shrinking X'_α without loss of generality

- (*)₃ if $\gamma \in X'_\alpha$ then $f_{\mu \times \alpha + \gamma}$ and $f_{\Upsilon_\varepsilon}^*$ are compatible.

For each $\gamma \in X'_\alpha$ let

$$t_\gamma = \{\beta \in X_{\Upsilon_\varepsilon} : f_{\mu \times \Upsilon_\varepsilon + \beta} \text{ and } f_{\mu \times \alpha + \gamma} \text{ are incompatible}\}.$$

As $\langle f_{\mu \times \Upsilon_\varepsilon + \beta} : \beta \in X_{\Upsilon_\varepsilon} \rangle$ is a Δ -system with heart $f_{\Upsilon_\varepsilon}^*$ (and $(*)_3$) necessarily $(*)_4$ $\gamma \in X'_\alpha$ implies t_γ is countable.

For $\gamma \in X'_\alpha$ let

$$s_\gamma \stackrel{\text{def}}{=} \bigcup \{u : u \text{ is a finite subset of } X'_\alpha \text{ and } F_\alpha(\{\mu \times \alpha + \beta : \beta \in u\}) \text{ belongs to } t_\gamma\}.$$

As F_α is a one-to-one function clearly

$(*)_5$ s_γ is a countable set.

Hence without loss of generality (possibly shrinking X'_α), as $\mu > \aleph_1$,

$(*)_6$ if $\gamma_1 \neq \gamma_2$ are from X'_α then $\gamma_1 \notin s_{\gamma_2}$.

By the choice of F_α for some finite subset u of X'_α with at least two elements, letting $u' \stackrel{\text{def}}{=} \{\mu \times \alpha + j : j \in u\}$ we have

$$\beta \stackrel{\text{def}}{=} F_\alpha(u') \in \{\mu \times \gamma(g_\varepsilon, \zeta_\varepsilon) + \gamma : \gamma \in X_{\gamma(g_\varepsilon, \zeta_\varepsilon)}\}$$

(remember $\Upsilon_\varepsilon = \gamma(g_\varepsilon, \zeta_\varepsilon)$), so $u' \cup \{\beta\} \in W$. Thus it is enough to show that $\{f_{\mu \times \alpha + j} : j \in u\} \cup \{f_\beta\}$ are compatible. For this it is enough to check any two. Now, $\{f_{\mu \times \alpha + j} : j \in u\}$ are compatible as $\langle f_{\mu \times \alpha + j} : j \in X_\alpha \rangle$ is a Δ -system. So let $j \in u$, why are $f_{\mu \times \alpha + j}$, f_β compatible? As otherwise $\beta - (\mu \times \Upsilon_\varepsilon) \in t_j$ and hence u is a subset of s_j . But u has at least two elements, so there is $\gamma \in u \setminus \{j\}$. Now u is a subset of X'_α and this contradicts the statement $(*)_6$ above, finishing the proof. $\blacksquare_{1.7}$

Remark 1.8. In 1.7, we can also get $d(BA(W, \mathbf{w})) = \mu$, but this is irrelevant to our aim. E.g. in this case let for $i < \mu$, h_i be a partial function from 2^μ to $\{0, 1\}$ such that $\text{Dom}(h_i) \cap [\beta, \beta + \mu)$ is finite for $\beta < 2^\mu$ and such that every finite such function is included in some h_i . Choosing the $(W_\alpha, \mathbf{w}_\alpha)$ preserve:

$$\{x_\beta : h_i(\beta) = 1\} \cup \{-x_\beta : h_i(\beta) = 0\} \text{ generates a filter of } BA(W_\alpha, \mathbf{w}_\alpha).$$

Conclusion 1.9. Theorem 0.1 holds.

PROOF By 1.1, 1.7. $\blacksquare_{2.1}$

2. Getting the example for $\mu = (\aleph_2)^{\aleph_0}$, $\lambda = 2^{\aleph_2}$. Our aim here is to show that there are $I, \mathfrak{B}$ as in 0.1 for $\mu = (\aleph_2)^{\aleph_0}$. For this we shall weaken the conditions in the Main Lemma 1.1 (see 2.1 below) and then show that we can get it in a variant of 1.7 (see 2.2 below). More fully, by 2.2 there is a $2^{\aleph_2}$ -candidate $(W, \mathbf{w})$ satisfying the assumptions of 2.1 except possibly clause **(a)**, so μ is irrelevant in the clauses **(b)**–**(f)**. Let $\mu = (\aleph_2)^{\aleph_0} = \aleph_2 + 2^{\aleph_0}$ and apply 2.2. Now we get the conclusion of 1.1 as required.

Proposition 2.1. *Assume that*

- (a) $\mu = \mu^{\aleph_0}$, $\lambda \leq 2^\mu$,
- (b) $\mathfrak{B}$ is a complete c.c.c. Boolean Algebra,
- (c) $x_i \in \mathfrak{B} \setminus \{0\}$ for $i < \lambda$, and $\mathcal{S} \subseteq \{u \in [\lambda]^{\leq \aleph_0} : (\forall i \in \lambda \setminus u)(x_i \notin \mathfrak{B}_u)\}$, where $\mathfrak{B}_u$ is the completion of $\langle \{x_i : i \in u\} \rangle_{\mathfrak{B}}$ in $\mathfrak{B}$ (for $u \in [\lambda]^{\leq \aleph_0}$),
- (d)[−] if $i \in u_i \in [\lambda]^{\leq \aleph_0}$ for $i < \lambda$, then we can find $n < \omega$, $i_0 < \dots < i_{n-1} < \lambda$ and $u \in \mathcal{S}(\subseteq [\lambda]^{\leq \aleph_0})$ such that:
 - (i) $\mathfrak{B} \models \bigcap_{\ell < n} x_{i_\ell} = 0$,
 - (ii) $i_\ell \in u_{i_\ell} \setminus u$ for $\ell < n$,
 - (iii) $\langle u_{i_\ell} \setminus u : \ell < n \rangle$ are pairwise disjoint;
- (e) $u \in \mathcal{S} \ \& \ i \in \lambda \setminus u \ \& \ y \in \mathfrak{B}_u \setminus \{0, 1\} \Rightarrow y \cap x_i \neq 0 \ \& \ y - x_i \neq 0$,
- (f) $\mathcal{S}$ is cofinal in $([\mu]^{< \aleph_0}, \subseteq)$
[actually, it follows from (d)[−]].

Then there are a σ -ideal I on $\mathcal{P}(\mu)$ and a σ -algebra $\mathfrak{A}$ of subsets of μ extending I such that $\mathfrak{A}/I$ satisfies the c.c.c. and the natural homomorphism $\mathfrak{A} \longrightarrow \mathfrak{A}/I$ cannot be lifted.

Remark: Actually we can in clause **(e)** omit “ $y - x_i \neq 0$ ”.

PROOF Repeat the proof of 1.1 till the definition of $\mathbf{e}_{i,n}$ and W_i in the beginning of the proof of 1.1.3 (which says that h_2 cannot be lifted). Then choose $u_i \in \mathcal{S}$ such that $W_i \subseteq \mathfrak{B}_{u_i}$ (exists by clause **(f)** of our assumptions). By clause **(d)**[−] we can find $n < \omega$, $i_0 < \dots < i_{n-1}$ and $u \in \mathcal{S}$ such that clauses **(i)**, **(ii)**, **(iii)** of **(d)**[−] hold.

Claim 2.1.1. *For $\ell < n$, there are homomorphisms f_{i_ℓ} from $\mathfrak{B}$ into $\{0, 1\}$ respecting $\mathbf{e}_{i_\ell, m}$ for $m < \omega$ and mapping x_{i_ℓ} to 1 such that $\langle f_{i_\ell} \upharpoonright (W_{i_\ell} \cap \mathfrak{B}_u) : \ell < n \rangle$ are compatible functions.*

Proof of the claim: E.g. by absoluteness it suffices to find it in some generic extension. Let $G_u \subseteq \mathfrak{B}_u$ be a generic ultrafilter. Now $\mathfrak{B}_u \triangleleft \mathfrak{B}$ and $(\forall y \in G_u)(y \cap x_{i_\ell} > 0)$ (see clause **(e)**). So there is a generic ultrafilter G_ℓ of $\mathfrak{B}$ extending G_u such that $x_{i_\ell} \in G_\ell$. Define f_{i_ℓ} by $f_{i_\ell}(y) = 1 \Leftrightarrow y \in G_\ell$

for $y \in u_{i_\ell}$. By Clause (iii) of $(\mathbf{d})^-$ those functions are compatible and we finish as in 1.1.

Thus we have finished. ■_{2.1}

Theorem 2.2. *In 1.7 if we let e.g. $\mu = \aleph_2$ then we can find a 2^μ -candidate $(W, \mathbf{w})$ such that $BA^c(W, \mathbf{w})$ satisfies the clauses (b)–(f) of 2.1.*

PROOF In short, we repeat the proof of 1.7 after defining $(W, \mathbf{w})$. But now we are being given $\langle u_i : i < \lambda \rangle$, $u_i \in [2^\mu]^{\leq \aleph_0}$, $i \in u_i$. For each $\alpha < 2^\mu$ (we cannot in general find a Δ -system but) we can find u_α^* , X_α such that $X_\alpha \in [\mu]^\mu$, $u_\alpha^* \in \mathcal{S} \subseteq [2^\mu]^{\leq \aleph_0}$ and $\langle u_{\mu \times \alpha + i} \setminus u_\alpha^* : i \in X_\alpha \rangle$ are pairwise disjoint, and $i \in X_\alpha \Rightarrow \mu \times \alpha + i \in u_{\mu \times \alpha + i} \setminus u_\alpha^*$ and we continue as there (replacing the functions by the sets where instead $G_\zeta = \{g : g \in Z_\zeta, \text{Dom}(g) \subseteq Z_\zeta\}$ we let h_ζ be a one-to-one function from Z_ζ onto μ and $G_\zeta = \{u \subseteq Z_\zeta : h_\zeta''(u) \in \mathcal{S}\}$ and instead $g = f_\alpha^* \upharpoonright Z_{\omega_1}$ let $u_\alpha^* \cap Z_{\omega_1} \subseteq Z_{\zeta_0(*)}$, $u_\alpha^* \cap Z_{\omega_1} \subseteq v \in G_\zeta$).

DETAILED PROOF Let $F^* : [\mu]^{< \aleph_0} \longrightarrow \mu$ be such that

$$(\forall A \in [\mu]^\mu)[F''([A]^{< \aleph_0} \setminus [A]^{< 2}) = \mu].$$

Let $\langle \bar{A}^\alpha : \alpha < 2^\mu \rangle$ list the sequences $\bar{A} = \langle A_i : i < \mu \rangle$ such that $A_i \in [2^\mu]^\mu$, $(\forall i < \mu)(\exists \alpha)(A_i \subseteq [\mu \times \alpha, \mu \times \alpha + \mu))$ and $i < j < \mu \Rightarrow A_i \cap A_j = \emptyset$. Without loss of generality we have $A_i^\alpha \subseteq \mu \times (1 + \alpha)$ and each $\bar{A}$ is equal to $\bar{A}^\alpha$ for 2^μ ordinals α . Clearly $\text{otp}(A_i^\alpha) = \mu$.

We choose by induction on $\alpha < 2^\mu$ pairs $(W_\alpha, \mathbf{w}_\alpha)$ and functions F_α such that

- (α) $(W_\alpha, \mathbf{w}_\alpha)$ is a $\mu \times (1 + \alpha)$ -candidate,
- (β) $\beta < \alpha$ implies $W_\beta = W_\alpha \cap [\mu \times (1 + \beta)]^{< \aleph_0}$, $\mathbf{w}_\beta = \mathbf{w}_\alpha \upharpoonright W_\beta$,
- (γ) F_α is a one-to-one function from

$\{u : u \subseteq [\mu \times (1 + \alpha), \mu \times (1 + \alpha + 1)) \text{ finite with at least two elements}\}$

into $\bigcup_{i < \mu} A_i^\alpha$,

- (δ) $W_{\alpha+1} = W_\alpha \cup \{u \cup \{F_\alpha(u)\} : u \in W_\alpha^*\}$, where $W_\alpha^* = \{u : u \text{ is a subset of } [\mu \times (1 + \alpha), \mu \times (1 + \alpha + 1)) \text{ such that } \aleph_0 > |u| \geq 2\}$,
- (ε) for finite $u \in W_\alpha^*$ we have

$$\mathbf{w}(u \cup \{F_\alpha(u)\}) = \{v \subseteq u \cup \{F_\alpha(u)\} : u \subseteq v \text{ or } F_\alpha(u) \in v \text{ \& } v \cap u \neq \emptyset\},$$

- (ζ) Let F_α be such that for any subset X of $J_\alpha = [\mu \times (1 + \alpha), \mu \times (1 + \alpha + 1))$ of cardinality μ and $i < \mu$ and $\gamma \in A_i^\alpha$ for some finite subset u of X we have $F_\alpha(u) \in A_i^\alpha \setminus \gamma$.

There are no difficulties in carrying out the construction and checking that it as required. Let $W = \bigcup_{\alpha} W_{\alpha}$, $\mathbf{w} = \bigcup_{\alpha} \mathbf{w}_{\alpha}$, $\mathfrak{B} = BA^c(W, \mathbf{w})$. Clearly $(W, \mathbf{w})$ is a λ -candidate.

Let $\mathcal{S}^* \subseteq [\mu]^{\leq \aleph_0}$ be stationary of cardinality μ . Let

$$\mathcal{S}' = \{u \in [\lambda]^{\leq \aleph_0} : \text{if } v \in W \text{ and } v \cap u \in \mathbf{w}(v) \text{ then } v \subseteq u\}.$$

Now, clause (f) holds as $(W, \mathbf{w})$ satisfies clause (d) of Definition 1.3(3). As for clause (e) use Lemma 2.3 below.

The main point is clause **(d)**⁻ of 2.1. So let $i \in a_i \in [\lambda^{\mu}]^{\leq \aleph_0}$ for $i < \lambda$ be given. For each $\alpha < \lambda$, as $\mu = \aleph_2$ we can find $X_{\alpha} \in [\mu]^{\mu}$ and $a_{\alpha}^* \in \mathcal{S}'$ such that $\alpha \in a_{\alpha}^*$ and:

$$\begin{aligned} (\otimes_{\alpha}) \quad \zeta_1 \neq \zeta_2 \ \& \ \zeta_1 \in X_{\alpha} \ \& \ \zeta_2 \in X_{\alpha} \quad \Rightarrow \quad a_{\mu \times \alpha + \zeta_1} \cap a_{\mu \times \alpha + \zeta_2} \subseteq a_{\alpha}^* \text{ and} \\ \zeta \in X_{\alpha} \quad \Rightarrow \quad \mu \times \alpha + \zeta \notin a_{\alpha}^*. \end{aligned}$$

For each $b \in [\lambda]^{\leq \aleph_0}$ let $\langle \gamma(b, i) : i < i(b) \rangle$ be a maximal sequence such that $\gamma(b, i) < \lambda$ and $u_{\gamma(b, i)}^* \cap u_{\gamma(b, j)}^* \subseteq b$ and $\gamma(b, i) \notin b$ for $j < i$ (just choose $\gamma(b, i)$ by induction on i).

We choose by induction on $\zeta \leq \omega_1$, $Y_{\zeta}, h_{\zeta}, S_{\zeta}, G_{\zeta}, Z_{\zeta}$ and $U_{\zeta, g}$ such that

- (a) $Y_{\zeta} \in [2^{\mu}]^{\leq \mu}$ is increasing continuous in ζ ,
- (b) Z_{ζ} is the minimal subset of λ (of cardinality $\leq \mu$) which includes

$$\bigcup \{u_{\gamma} : (\exists \alpha \in Y_{\zeta}) [\mu \times \alpha \leq \gamma < \mu \times (\alpha + 1)]\}$$

and satisfies

$$u \in W \ \& \ u \cap Z_{\zeta} \in \mathbf{w}(u) \quad \Rightarrow \quad u \subseteq Z_{\zeta},$$

- (c) h_{ζ} is a one-to-one function from μ onto Z_{ζ} , and

$$G_{\zeta} = \{h_{\zeta}''(b) : b \in \mathcal{S}\} \cup \bigcup_{\xi < \zeta} G_{\xi}.$$

- (d) for $b \in G_{\zeta}$ we have $U_{\zeta, b}$ is $\{i : i < i(b)\}$ if $i(b) < \mu^+$ and otherwise is a subset of $i(b)$ of cardinality μ such that

$$j \in U_{\zeta, b} \quad \Rightarrow \quad \text{Dom}(f_{\gamma(b, j)}^*) \cap Z_{\zeta} \subseteq b,$$

- (e) $Y_{\zeta+1} = Y_{\zeta} \cup \{\gamma(b, j) : b \in G_{\zeta} \text{ and } j \in U_{\zeta, b}\}.$

Again, there is no problem to carry out the definition (e.g. $|Z_{\zeta}| \leq \mu$ by clause (d) of 1.3(3)). Let $Y = Y_{\omega_1}$. Let $\{(b_{\varepsilon}, \xi_{\varepsilon}) : \varepsilon < \varepsilon(*) \leq \mu\}$ list the set of pairs (b, ξ) such that $\xi < \omega_1$, $b \in G_{\xi}$ and $i(b) \geq \mu^+$. We can find $\langle \zeta_{\varepsilon} : \varepsilon < \varepsilon(*) \rangle$ such that $\langle \gamma(b_{\varepsilon}, \zeta_{\varepsilon}) : \varepsilon < \varepsilon(*) \rangle$ is without repetition and $\zeta_{\varepsilon} \in U_{b_{\varepsilon}, \xi_{\varepsilon}}$, $\varepsilon(*) \leq \mu$. So for some $\alpha < 2^{\mu} \setminus Y_{\omega_1}$ we have

$$(\forall \varepsilon < \varepsilon(*))(A_{\varepsilon}^{\alpha} = \{\mu \times \gamma(b_{\varepsilon}, \zeta_{\varepsilon}) + \Upsilon : \Upsilon \in X_{\gamma(b_{\varepsilon}, \zeta_{\varepsilon})}\}.$$

Now, let $b_0 = a_\alpha^* \cap Z_{\omega_1}$, so for some $\zeta_0(*) < \omega_1$ we have $b_0 \subseteq Z_{\zeta_0(*)}$. As a_α^* is countable and $G_\zeta \subseteq [Z_\zeta]^{\leq \aleph_0}$ is stationary (and the closure property of Z_ζ) there is $b^* \in \mathcal{S}'$ such that $b \stackrel{\text{def}}{=} b^* \cap Z_{\zeta_0(*)}$ belongs to G_ζ and $a_\alpha^* \subseteq b^*$ and so $U_{b,\zeta} \subseteq i(b)$ for $\zeta \in [\zeta_0(*), \omega_1)$ and $\langle \gamma(b, i) : i < i(b) \rangle$ are well defined. Now α exemplified $i(b) < \mu^+$ is impossible (see the maximality as otherwise $i < i(b) \Rightarrow \gamma(b, i) \in Z_{\zeta_0(*)+1} \subseteq Z_{\omega_1}$).

As for each $\gamma \in X_\alpha$, the set $a_{\mu \times \alpha + \gamma}$ is countable, for some $\zeta_{1,\gamma}(*) < \omega_1$ we have $a_{\mu \times \alpha + \gamma} \cap Z_{\omega_1} \subseteq Z_{\zeta_{1,\gamma}(*)}$. Since $\text{cf}(\mu) > \aleph_1$ necessarily for some $\zeta_1(*) < \omega_1$ we have

$$X'_\alpha \stackrel{\text{def}}{=} \{\gamma \in X_\alpha : \zeta_{1,\gamma}(*) \leq \zeta_1(*)\} \in [\mu]^\mu$$

and without loss of generality $\zeta_1(*) \geq \zeta_0(*)$. Thus for some $\varepsilon < \mu$ we have $b_\varepsilon = b \ \& \ \xi_\varepsilon = \zeta_1(*) + 1$. Let $\Upsilon_\varepsilon = \gamma(b_\varepsilon, \zeta_\varepsilon)$. Clearly

- (*)₁ $a_\alpha^*, a_{\Upsilon_\varepsilon}^*$ are countable,
- (*)₂ $\gamma \in X'_\alpha \Rightarrow \mu \times \alpha + \gamma \in a_\gamma$,
- (*)₃ $\gamma_1 \neq \gamma_2 \ \& \ \gamma_1 \in X'_\alpha \ \& \ \gamma_2 \in X'_\alpha \Rightarrow a_{\mu \times \alpha + \gamma_1} \cap a_{\mu \times \alpha + \gamma_2} \subseteq b^*$.

So possibly shrinking X'_α without loss of generality

- (*)₄ if $\gamma \in X'_\alpha$ then $a_{(\mu \times \alpha + \gamma)} \cap a_{\Upsilon_\varepsilon}^* \subseteq b^*$.

For each $\gamma \in X'_\alpha$ let

$$t_\gamma = \{\beta \in X_{\Upsilon_\varepsilon} : a_{(\mu \times \Upsilon_\varepsilon + \beta)} \cap a_{(\mu \alpha + \gamma)} \not\subseteq b^*\}.$$

As $\langle f_{(\mu \times \Upsilon_\varepsilon + \beta)} : \beta \in X_{\Upsilon_\varepsilon} \rangle$ was chosen to satisfy $(\otimes_{\Upsilon_\varepsilon})$ (and (*)₃) necessarily

- (*)₅ $\gamma \in X'_\alpha$ implies t_γ is countable.

For $\gamma \in X'_\alpha$ let

$$s_\gamma \stackrel{\text{def}}{=} \bigcup \{u : u \text{ is a finite subset of } X'_\alpha \text{ and } F_\alpha(\{\mu \times \alpha + \beta : \beta \in u\}) \text{ belongs to } t_\gamma\}.$$

As F_α is a one-to-one function clearly

- (*)₆ s_γ is a countable set.

So without loss of generality (possibly shrinking X'_α using $\mu > \aleph_1$)

- (*)₇ if $\gamma_1 \neq \gamma_2$ are from X'_α then $\gamma_1 \notin s_{\gamma_2}$.

By the choice of F_α , for some finite subset u of X'_α with at least two elements, letting $u' \stackrel{\text{def}}{=} \{\mu \times \alpha + j : j \in u\}$ we have

$$\beta \stackrel{\text{def}}{=} F_\alpha(u') \in \{\mu \times \gamma(b_\varepsilon, \zeta_\varepsilon) + \gamma : \gamma \in X_{\gamma(b_\varepsilon, \zeta_\varepsilon)}\}.$$

Hence $u' \cup \{\beta\} \in W$, so it is enough to show that $\{a_{\mu \times \alpha + j} : j \in u\} \cup \{a_\beta\}$ are pairwise disjoint outside b^* . For the first it is enough to check any two. Now, $\{f_{\mu \times \alpha + j} : j \in u\}$ are O.K. by the choice of $\langle f_{\mu \times \alpha + j} : j \in X_\alpha \rangle$. So let

$j \in u$. Now, $a_{\mu \times \alpha + j}$, a_β are O.K., otherwise $\beta - (\mu \times \Upsilon_\varepsilon) \in t_j$ and hence u is a subset of s_j but u has at least two elements and is a subset of X'_α and this contradicts the statement $(*)_6$ above and so we are done. $\blacksquare_{2.2}$

Lemma 2.3. *Let $(W, \mathbf{w})$ be a λ -candidate. Assume that $u \subseteq \lambda$ and $u = \text{cl}_{(W, \mathbf{w})}(u)$ (see Definition 1.3(1),(d)) and let $W^{[u]} = W \cap [u]^{<\aleph_0}$ and $\mathbf{w}^{[u]} = \mathbf{w} \upharpoonright W^{[u]}$. Furthermore suppose that $(W, \mathbf{w})$ is non-trivial (which holds in all the cases we construct), i.e.*

$$(*) \quad i \in v \in W \quad \Rightarrow \quad v \setminus \{i\} \in \mathbf{w}(v).$$

Then:

1. $(W^{[u]}, \mathbf{w}^{[u]})$ is a λ -candidate (here $u = \text{cl}_{(W, \mathbf{w})}(u)$ is irrelevant);
2. $BA(W^{[u]}, \mathbf{w}^{[u]})$ is a subalgebra of $BA(W, \mathbf{w})$, moreover $BA(W^{[u]}, \mathbf{w}^{[u]}) \triangleleft BA(W, \mathbf{w})$;
3. if $i \in \lambda \setminus u$ and $y \in BA(W^{[u]}, \mathbf{w}^{[u]})$ then
$$y \neq 0 \quad \Rightarrow \quad y \cap x_i > 0 \ \& \ y - x_i > 0;$$
4. $BA^c(W^{[u]}, \mathbf{w}^{[u]}) \triangleleft BA^c(W, \mathbf{w})$.

PROOF 1) Trivial.

2) *The first phrase:* if f_0 is a homomorphism from $BA(W^{[u]}, \mathbf{w}^{[u]})$ to the Boolean Algebra $\{0, 1\}$ we define a function f from $\{x_\alpha : \alpha < \lambda\}$ to $\{0, 1\}$ by $f(x_\alpha)$ is $f_0(x_\alpha)$ if $\alpha \in u$ and is zero otherwise. Now

$$v \in W \quad \Rightarrow \quad (\exists \alpha \in v)(f(x_\alpha) = 0).$$

Why? If $v \subseteq u$, then $v \in W^{[u]}$ and “ f_0 is a homomorphism”, so $f_0(\bigcap_{\alpha \in v} x_\alpha) = 0$. Hence $(\exists \alpha \in v)(f_0(x_\alpha) = 0)$ and hence $(\exists \alpha \in v)(f(x_\alpha) = 0)$. If $v \not\subseteq u$, then choose $\alpha \in v \setminus u$, so $f(x_\alpha) = 0$.

So f respects all the equations involved in the definition of $BA(W, \mathbf{w})$ hence can be extended to a homomorphism $\hat{f}$ from $BA(W, \mathbf{w})$ to $\{0, 1\}$. Easily $f_0 \subseteq \hat{f}$ and so we are done.

As for *the second phrase*, let $z \in BA(W, \mathbf{w})$, $z > 0$ and we shall find $y \in BA(W^{[u]}, \mathbf{w}^{[u]})$, $y > 0$ such that

$$(\forall x)[x \in BA(W^{[u]}, \mathbf{w}^{[u]}) \ \& \ 0 < x \leq y \quad \Rightarrow \quad x \cap z \neq 0].$$

We can find disjoint finite subsets s_0, s_1 of λ such that $0 < z' \leq z$ where $z' = \bigcap_{\alpha \in s_1} x_\alpha \cap \bigcap_{\alpha \in s_0} (-x_\alpha)$. Let

$$t = \bigcup \{v : v \in W \text{ a finite subset of } \lambda \text{ and } v \cap s_0 \in \mathbf{w}(v)\} \cup s_0 \cup s_1.$$

We know that t is finite. We can find a partition t_0, t_1 of t (so $t_0 \cap t_1 = \emptyset$, $t_0 \cup t_1 = t$) such that $s_0 \subseteq t_0$ and $s_1 \subseteq t_1$ and $y^* = \bigcap_{\alpha \in t_1} x_\alpha \cap \bigcap_{\alpha \in t_0} (-x_\alpha) >$

0. Note that $y \stackrel{\text{def}}{=} \bigcap_{\alpha \in u \cap t_1} x_\alpha \cap \bigcap_{\alpha \in u \cap t_0} (-x_\alpha)$ is > 0 and, of course, $y \in BA(W^{[u]}, \mathbf{w}^{[u]})$. We shall show that y is as required. So assume $0 < x \leq y$, $x \in BA(W^{[u]}, \mathbf{w}^{[u]})$. As we can shrink x , without loss of generality, for some disjoint finite $r_0, r_1 \subseteq u$ we have $t \cap u \subseteq r_0 \cup r_1$ and $x = \bigcap_{\alpha \in r_1} x_\alpha \cap \bigcap_{\alpha \in r_0} (-x_\alpha)$, so clearly $t_1 \cap u \subseteq r_1$, $t_0 \cap u \subseteq r_0$.

We need to show $x \cap z \neq 0$, and for this it is enough to show that $x \cap z' \neq 0$. Now, it is enough to find a function $f : \{x_\alpha : \alpha < \lambda\} \rightarrow \{0, 1\}$ respecting all the equations in the definition of $BA(W, \mathbf{w})$ such that $\hat{f}$ maps $x \cap z'$ to 1. So let $f(x_\alpha) = 1$ for $\alpha \in r_1 \cup s_1$ and $f(x_\alpha) = 0$ otherwise. If this is O.K., fine as $f \upharpoonright r_0, f \upharpoonright s_0$ are identically zero and $f \upharpoonright r_1, f \upharpoonright s_1$ are identically one. If this fails, then for some $v \in \mathbf{w}$ we have $v \subseteq r_1 \cup s_1$. But then $v \cap r_1 \in \mathbf{w}(v)$ or $v \cap s_1 \in \mathbf{w}(v)$. Now if $v \cap r_1 \in \mathbf{w}(w)$ as $r_1 \subseteq u$ necessarily $v \subseteq u$, but $v \subseteq r_1 \cup s_1$ and $s_1 \cap u \subseteq t_1 \subseteq r_1$, so $v \subseteq r_1$ is a contradiction to $x > 0$. Lastly, if $v \cap s_1 \in \mathbf{w}(v)$, then $v \subseteq t$ so as $v \subseteq r_1 \cup s_1$ we have $v \subseteq s_1 \cup (t \cap r_1)$ and so $v \subseteq s_1 \cup t_1$ and hence $v \subseteq t_1$ — a contradiction to $y^* > 0$. So f is O.K. and we are done.

3) Let f_0 be a homomorphism from $BA(W^{[u]}, \mathbf{w}^{[u]})$ to the trivial Boolean Algebra $\{0, 1\}$. For $t \in \{0, 1\}$ we define a function f from $\{x_\alpha : \alpha < \lambda\}$ to $\{0, 1\}$ by

$$f(x_\alpha) = \begin{cases} f_0(x_\alpha) & \text{if } \alpha \in u \\ t & \text{if } \alpha = i \\ 0 & \text{if } \alpha \in \lambda \setminus u \setminus \{i\}. \end{cases}$$

Now f respects the equations in the definition of $BA(W, \mathbf{w})$. Why? Let $v \in W$. We should prove that $(\exists \alpha \in v)(f(\alpha) = 0)$. If $v \subseteq u$, then

$$f \upharpoonright \{x_\alpha : \alpha \in v\} = f_0 \upharpoonright \{x_\alpha : \alpha \in v\} \quad \text{and}$$

$$0 = f_0(0_{BA(W^{[u]}, \mathbf{w}^{[u]})}) = f_0\left(\bigcap_{\alpha \in v} x_\alpha\right) = \bigcap_{\alpha \in v} f_0(x_\alpha),$$

so $(\exists \alpha \in v)(f_0(x_\alpha) = 0)$. If $v \not\subseteq u \cup \{i\}$ let $\alpha \in v \setminus u \setminus \{i\}$, so $f(x_\alpha) = 0$ as required.

So we are left with the case $v \subseteq u \cup \{i\}$, $v \not\subseteq u$. Then by the assumption (*), $v \cap u = v \setminus \{i\} \in \mathbf{w}(v)$ and $v \subseteq u$, a contradiction.

4) Follows. ■_{2.3}

Remark 2.4. We can replace $\aleph_0$ by say $\kappa = \text{cf}(\kappa)$ (so in 2.2, $\mu = \kappa^{++}$, in 1.7, $(\forall \alpha < \mu)(|\alpha|^{<\kappa} < \mu = \text{cf}(\mu))$).

REFERENCES

- [1] ENGELKING, R. and KARLOWICZ, M., *Some theorems of set theory and their topological consequences*, Fundamenta Math., 57 (1965), 275–285.
- [2] FREMLIN, D., *Measure algebras*, In *Handbook of Boolean Algebras*, North-Holland, 1989. Monk D., Bonnet R. eds.
- [3] DETLEFSEN, M. E. and SZYMAŃSKI, A., *Category bases*, International Journal of mathematics and Mathematical Sciences, 16 (1993), 531–538.
- [4] GITIK, M. and SHELAH, S., *On densities of box products*, Topology and its Applications, accepted.
- [5] HAJNAL, A. and JUHASZ, I. and SZENTMIKLOSSY, Z., *On the structure of CCC partial orders*, Algebra Universalis, to appear.
- [6] MORGAN, J. C. II, *Point set theory*, volume 131 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker, Inc, New York, 1990.
- [7] RABUS, M. and SHELAH, S., *Topological density of ccc Boolean algebras - every cardinality occurs*, Proceedings of the American Mathematical Society, submitted.
- [8] SHELAH, S., *Remarks on λ -collectionwise Hausdorff spaces*, Topology Proceedings, 2 (1977), 583–592.
- [9] SHELAH, S., *On saturation for a predicate*. Notre Dame Journal of Formal Logic, 22 (1981), 239–248.
- [10] SHELAH, S., *Cardinal Arithmetic*, volume 29 of *Oxford Logic Guides*. Oxford University Press, 1994.
- [11] SHILLING, K., *Some category bases which are equivalent to topologies*, Real Analysis Exchange, 14 (1988/89), 210–214.
- [12] SZYMAŃSKI, A., *On the measurability problem for category bases*, Real Analysis Exchange, 17 (1991/92), 85–92.

INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY, JERUSALEM 91904, ISRAEL,
AND RUTGERS UNIVERSITY, MATHEMATICS DEPARTMENT, NEW BRUNSWICK, NJ 08854,
USA

E-mail address: `shelah@math.huji.ac.il`